

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NUSC TD 7983			7a. NAME OF MONITORING ORGANIZATION	
6a. NAME OF PERFORMING ORGANIZATION Naval Underwater Systems Center		6b. OFFICE SYMBOL (if applicable) 10		7b. ADDRESS (City, State, and ZIP Code)
6c. ADDRESS (City, State, and ZIP Code) New London Laboratory New London, CT 06320			9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Naval Underwater Systems Center		8b. OFFICE SYMBOL (if applicable) 10		10. SOURCE OF FUNDING NUMBERS
8c. ADDRESS (City, State, and ZIP Code) New London Laboratory New London, CT 06320			PROGRAM ELEMENT NO.	PROJECT NO. 710Y11
			TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) RESOLUTION OF LOW-FREQUENCY ATTENUATION ANOMALIES IN THE SOUTHERN HEMISPHERE				
12. PERSONAL AUTHOR(S) R. N. Denham, R. W. Bannister, K. M. Guthrie, R. H. Mellen, and D. G. Browning				
13a. TYPE OF REPORT		13b. TIME COVERED FROM TO		15. PAGE COUNT 12
14. DATE OF REPORT (Year, Month, Day) 1987 June 1				
16. SUPPLEMENTARY NOTATION A paper presented at the 113th meeting of the Acoustical Society of America, 11-15 May 1987, Indianapolis, Indiana				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	Sound Absorption	
			Sound Attenuation	
			South Pacific Ocean	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) During Project Kiwi One (D. G. Browning et al., Nature, 282, 820-822 (1979)) anomalously high values of low-frequency attenuation were observed in the central South Pacific Ocean. Similar results were obtained along a track from New Zealand into the Southern Ocean during Project Tasman Two (R. W. Bannister et al., J. Acoust. Soc. Am., 62, 847-859 (1977)). A recent analysis of oceanographic data by Mellen shows that both regions are included in a relatively high pH contour at the sound channel axis. The corresponding predicted values of attenuation are in reasonable agreement with the measured data.				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL David G. Browning			22b. TELEPHONE (Include Area Code) (203) 440-4173	22c. OFFICE SYMBOL 3331

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted.
All other editions are obsolete.SECURITY CLASSIFICATION OF THIS PAGE
UNCLASSIFIED

NUSC Technical Document 7983
1 June 1987

Resolution of Low-Frequency Attenuation Anomalies in the Southern Hemisphere

**A Paper Presented at the
113th Meeting of the Acoustical Society of America,
11-15 May 1987, Indianapolis, Indiana**

R. N. Denham
R. W. Bannister
K. M. Guthrie
Defence Scientific Establishment

R. H. Mellen
PSI-Marine Sciences

D. G. Browning
Surface Ship Sonar Department




Naval Underwater Systems Center
/Newport, Rhode Island / New London, Connecticut

PREFACE

This document was prepared under NUSC
Project No. 710Y11.

REVIEWED AND APPROVED: 1 June 1987

A handwritten signature in dark ink, appearing to read 'W. A. Von Winkle'.

**W. A. VON WINKLE
ASSOCIATE TECHNICAL DIRECTOR FOR
RESEARCH AND TECHNOLOGY**



RESOLUTION OF LOW FREQUENCY ATTENUATION ANOMALIES IN THE SOUTHERN HEMISPHERE

**R. N. DENHAM, R. W. BANNISTER, K. M. GUTHRIE
DEFENCE SCIENTIFIC ESTABLISHMENT
NAVAL BASE POST OFFICE
AUCKLAND, NEW ZEALAND**


**R. H. MELLEN
PSI — MARINE SCIENCES
95 TRUMBULL STREET
NEW LONDON, CT 06320**

**D. G. BROWNING
NAVAL UNDERWATER SYSTEMS CENTER
NEW LONDON LABORATORY
NEW LONDON, CT 06320**

N0211-GA-87(L)-
00341.1 T


VIEWGRAPH 1

There are several areas of the South Pacific Ocean and adjacent Southern Ocean where the absorption component of low frequency attenuation appeared to be higher than predicted. An analysis of pH profiles in the southern hemisphere has resolved this dilemma. These results will be reported in this paper.



VIEWGRAPH 2

In the 1970's we conducted what we hoped would be the definitive attenuation experiment in the South Pacific Ocean: Project Kiwi One. The uniformity of the sound channel axis depth across the vast ocean allowed an extremely long data base of 10,000 km. We found, however, that the measured attenuation was not uniform but divided into three distinct regions which we termed west, central, and east.




VIEWGRAPH 3

The attenuation in the east and west regions was pretty much as expected. Data from the central region was significantly higher. It appeared related to the strong influx of intermediate water from the south but at the time we just had no good explanation.



PROJECT TASMAN TWO PROPAGATION EXPERIMENT




N0211-GA-87(LI)-
00341.4 T

VIEWGRAPH 4

Many of the southern hemisphere water masses run parallel to the dominant Antarctic circumpolar current. In the late 1970's we ran a series of acoustic tracks from the southern tip of New Zealand. These were designed to lay in and/or cross specific water masses. Only the southernmost track, designated as SW, produced anomalous results.



PROJECT TASMAN TWO ATTENUATION RESULTS




N0211-GA-87(LI-
00341.5 T

VIEWGRAPH 5

These anomalously high values of attenuation observed along the SW track were very similar to those measured in the central region of the Kiwi One track. It was not obvious, however, what the Southern Ocean and the central South Pacific had in common.



pH PROFILES SOUTH PACIFIC OCEAN



N0211-GA-87(L)-
00341.7 T

VIEWGRAPH 8


For the South Pacific, typical pH profiles show the influence of the relatively low pH North Pacific down to 10° south latitude, then the influence of the Antarctic circulation is felt and the profiles become more uniform. However, if the sound channel axis becomes shallower at higher latitudes, the attenuation for SOFAR propagation becomes greater even though the pH profile hasn't, because we have moved up the curve to higher pH values.



PACIFIC OCEAN **ABSORPTION (K-FACTOR) CONTOURS** **AT SOUND CHANNEL AXIS**

K = 10 (pH-8)


TRACKS OF PROPAGATION EXPERIMENTS



N0211-GA-87(L)-
00341.9 T

VIEWGRAPH 9

If we plot absorption (k-factor) contours for the Pacific Ocean based on a recent analysis by Bob Mellen, we can see both the cause of the anomaly and the reason for the similarity of the Kiwi One and Tasman Two results. The highest absorption contour encompasses the central region of the Kiwi One track and the SW track of Tasman Two.



VIEWGRAPH 10

For frequencies of 200 Hz and above the data from both regions blend in well with the predicted absorption values. Below 200 Hz the attenuation levels off as a mechanism other than absorption begins to dominate. We believe this is due to scattering but no one has conclusively proved this yet.



CONCLUSIONS

- BOTH ANOMALOUS REGIONS ARE WITHIN THE SAME pH CONTOUR
- VALUES OF ABSORPTION ARE IN REASONABLE AGREEMENT
- CAUSE OF LOW FREQUENCY SCATTERING COMPONENT IS NOT YET DETERMINED

N0211-GA-87(L)-
00341.11 T


VIEWGRAPH 11

In summary:

- o Both regions of high attenuation fall within the same pH contour.
- o Above 200 Hz the data are in good agreement with the predicted absorption.
- o We believe we can now accurately predict absorption for any ocean, however, the frequency independent attenuation below 200 Hz is not yet fully defined or understood.

INITIAL DISTRIBUTION LIST

Addressee	No. of Copies
DIA	1
DTIC	1
DARPA	1
NPS	1
NWC	1



U229484